

## **General Infusion Therapy Order Form**

atient Name:		DOB:	Phone:		
ddress:		City:	State:	Zip:	
. For new patients, please	submit with form:				
<ul> <li>Copy of insurance car</li> </ul>	d • Patient demographics	<ul><li>History and physical</li></ul>	<ul> <li>Pertinent labs and t</li> </ul>	est results based on therapy	
Patient Information					
Male Female Heigh	_	-	NKDA Allergies:		
Is this the first dose? Yes	No, date of last infusion:	Next due:	Line type:	PIV PICC Port Other:	
Diagnosis and Clinical In ICD-10 (required):	formation Primary diagnosis:		Other information:	ouici.	
Prescription Information	1				
Medication	Medication: Strength/formulation:				
Dose / Frequency	Dose:	Freque	ncy:		
	Route of administration: Intrav	enous Subcutaneous	Other:		
Directions	<ul><li>☑ Prepare and infuse medication</li><li>☐ Other:</li></ul>	per manufacturer guidelii	nes, as tolerated and per Pror	nptCare Policy & Procedure	
Quantity / Refills	☐ Dispense 1 month supply / QS on all selected medications / Refill x 12 months ☐ Other:  Dispense all medical supplies necessary for infusion				
Additional Orders					
Premedications: Give 30 m	in prior to infusions (Note: if nothing	g is checked, no premedic	ations will be given)		
Adults (or patients weighir	ng >40kg):	Pediatrics (weigh	ing <40 kg): (may adjust with	weight changes)	
$\hfill \Box$ Diphenhydramine 25-50mg PO. Patient may decline.		☐ Diphenhydran	☐ Diphenhydramine 1mg/kg PO		
$\hfill \square$ Acetaminophen 325-650mg PO. Patient may decline.		☐ Acetaminophe	☐ Acetaminophen 15mg/kg PO		
☐ Methylprednisolone 40mg (OR mg) slow IV push equivalent corticosteroid, substitution if needed by pharmacy)		. , , , , , , , , , , , , , , , , , , ,	☐ Methylprednisolone 1 mg/kg (OR mg) slow IV push (or an equivalent corticosteroid, substitution if needed by pharmacy)		
☐ Other:		ı			
For infusions to be adminis	tered by RN:				
	oplicable): RN to start peripheral IV	or use existing CVC. RN to	administer catheter flushing	per PromptCare Policy & Procedur	
RN may instruct pation	ent to hydrate pre/post infusion and	d educate on taking OTC o	iphenhydramine and/or acet	aminophen per manufacturer dosii	
recommendations as	needed to prevent/treat post-infus	ion headache.			
RN to monitor patier and when to contact Other:	nt for at least 30 min post infusion (o physician	or per protocol specific to	therapy) and educate on pos	sible side effects, allergic reactions	
Standard anaphylaxis	atients requiring an EpiPen in hom kit to be dispensed and dosed per I IM (50 mg/mL vial), and NS IV. Addi	protocol if indicated (ther			
Prescriber Information					
Prescriber Name:		Office contact:			
		City:	Stat	te: Zip:	
Address:					
Address: Phone: License #:		Fax:	NPI:		

health plan to obtain any authorizations necessary to enable it to receive payment for services.

Confidential Health Information: Healthcare information is personal information related to a person's healthcare. It is being faxed to you after appropriate authorization and under circumstances that don't require authorization. You are obligated to maintain it in a safe, secure and confidential manner. Re-disclosure of this information is prohibited by law or appropriate customer/patient authorization is obtained. Unauthorized re-disclosure or failure to maintain confidentiality could subject you to penalties described in federal and state law. Important Warning: This message is intended for the use of the person or entity to who it is addressed and may contain information that is privileged and confidential, the disclosure of which is governed by applicable law. If the reader of this message is not the intended recipient, or the employee or agent responsible for delivering it to the intended recipient, you are hereby notified that any discrimination, distribution, or copying of this information is STRICTLY PROHIBITED. If you have received this message in error, please notify us immediately.