promptcare™
Specialty Pharmacy & Infusion Services
An AOM Infusion Company

Intake Specialist 1-866-776-6782
Fax: 800-815-6808
newreferral@promptcare.com

Krystexxa (pegloticase) | Order Form

Patient Name: _____ DOB: _____ Phone: _____
Address: _____ City: _____ State: _____ Zip: _____

1. For new patients, please submit with form:

☒ Copy of insurance card
☒ Patient demographics
☒ History & physical
☒ Labs/records including G6PD deficiency screening (if indicated*) and baseline uric acid levels

2. Patient Information

☐ Male ☐Female Height: _____ in/cm Weight: _____ lbs/kg Allergies: _____

Is this the first dose? ☐Yes ☐No, date of last infusion: _____ Line type: ☐ PIV ☐PICC ☐Port ☐Other

3. Diagnosis and Clinical Information

Primary diagnosis information: ☐ Gout ☐ Other: _____ ICD-10 (required): _____

Is there an immunomodulator prescribed? ☐ Yes ☐ No **If yes, please indicate**: ☐ Methotrexate ☐ Other _____

Is patient currently taking oral urate-lowering agents? ☐ No ☐ Yes . Oral urate-lowering agents should be discontinued prior to Krystexxa

G6PD deficiency screening and/or testing results:

☐ Patient evaluated by provider and is **not** at risk and will **not** be tested (or tested negative)
☐ Patient is at risk for G6PD deficiency, and test results are: ☐ Positive (contraindicated) ☐ Negative

Baseline serum uric acid level: _____mg/dL (Must be >6.0mg/dL to initiate Krystexxa)

- Uric acid levels are required to be drawn 1 to 2 days prior to each infusion, **referring provider to arrange lab draws locally**
  Name of lab facility: _____ Phone number for lab results: _____
- Please fax results to PromptCare at 800-815-6808 as soon as available
- A single uric acid of >6.0mg/dL will require follow-up with provider, but will not post-pone next infusion

4. Prescription Information

Medication	☒ **Krystexxa** (pegloticase)
Dosing / Frequency	☒ 8mg in 250mL sodium chloride 0.9% IV every 2 weeks
Administration	☒ Prepare and infuse per manufacturer guidelines. Infuse over no less than 2 hours and observe patient for at least 1 hour following infusion ☒ May infuse in patient home unless otherwise noted: _____
Quantity / Refills	Dispense 2-week supply on all selected medications; Refill x 12 months unless otherwise specified: _____ Dispense all medical supplies necessary for infusion

Confidential Health Information: Healthcare information is personal information related to a person's healthcare. It is being faxed to you after appropriate authorization and under circumstances that don't require authorization. You are obligated to maintain it in a safe, secure and confidential manner. Re-disclosure of this information is prohibited by law or appropriate customer/patient authorization is obtained. Unauthorized re-disclosure or failure to maintain confidentiality could subject you to penalties described in federal and state law. Important Warning: This message is intended for the use of the person or entity to who it is addressed and may contain information that is privileged and confidential, the disclosure of which is governed by applicable law. If the reader of this message is not the intended recipient, or the employee or agent responsible for delivering it to the intended recipient, you are hereby notified that any discrimination, distribution, or copying of this information is STRICTLY PROHIBITED. If you have received this message in error, please notify us immediately.

5. Additional Orders

☒ RN to start peripheral IV or use existing CVC. RN to administer catheter flushing per PromptCare Policy and Procedure

☒ **Give standard premedications 30 minutes prior to infusion:**

☒ Solu-medrol 40 mg **OR** _____ mg IV

☒ Acetaminophen 650mg PO

☒ Antihistamine: patient may take night prior AND morning of infusion. If nothing is checked, patient/pharmacist may select from the following OTC products:

☐ Diphenhydramine 25mg PO ☐ Cetirizine 10mg PO ☐ Fexofenadine 60 mg (Allegra 12 Hour) PO ☐ Fexofenadine 180 mg (Allegra 24 Hour) PO ☐ Loratadine 10mg PO

☐ Other: _____

☒ RN to instruct patient to hydrate pre/post infusion and educate on taking OTC diphenhydramine and/or acetaminophen per manufacturer dosing recommendations as needed to prevent/treat post-infusion headache.

☒ RN to monitor patient for at least 1 hour post infusion and educate on possible side effects, allergic reactions, and when to contact physician

6. Adverse Reaction Orders

Standard anaphylaxis kit to be dispensed and dosed per protocol: Epinephrine IM/SQ (1 mg/mL vial), diphenhydramine IV/IM (50 mg/mL vial), and NS IV. Additional orders: _____

7. Prescriber Information

Prescriber Name: _____ Office Contact: _____

Address: _____ City: _____ State: _____ Zip: _____

Phone: _____ Fax: _____ License No.: _____

DEA No.: _____ NPI: _____

_____ _____ _____ _____

Physician Signature (Substitution Permitted) Date **Physician Signature (Dispense as Written)** Date

Confidential Health Information: Healthcare information is personal information related to a person's healthcare. It is being faxed to you after appropriate authorization and under circumstances that don't require authorization. You are obligated to maintain it in a safe, secure and confidential manner. Re-disclosure of this information is prohibited by law or appropriate customer/patient authorization is obtained. Unauthorized re-disclosure or failure to maintain confidentiality could subject you to penalties described in federal and state law. Important Warning: This message is intended for the use of the person or entity to who it is addressed and may contain information that is privileged and confidential, the disclosure of which is governed by applicable law. If the reader of this message is not the intended recipient, or the employee or agent responsible for delivering it to the intended recipient, you are hereby notified that any discrimination, distribution, or copying of this information is STRICTLY PROHIBITED. If you have received this message in error, please notify us immediately.